arXiv:1603.00885v1 [math.CO] 2 Mar 2016

Contemporary Mathematics

The K_{n+5} and $K_{3^2,1^n}$ families are obstructions to n-apex.

Thomas W. Mattman and Mike Pierce

Abstract. We give evidence in support of a conjecture that the Y∇Y families of K_{n+5} and $K_{3^2,1^n}$ are minor minimal obstructions for the n-apex property

Introduction

A graph is **n-apex** if deleting n or fewer vertices results in a planar graph. Kuratowski [**K**] showed the 0-apex or **planar** graphs are characterized by the obstruction set $\mathcal{O}_0 = \{K_5, K_{3,3}\}$. Using the formulation of Wagner [**W**], a graph is planar if and only if it has neither K_5 nor $K_{3,3}$ as a minor. Recall that H is a **minor** of the graph G if H can be obtained by contracting edges in a subgraph of G. As the n-apex property is minor closed, it follows from the Graph Minor Theorem of Robertson and Seymour [**RS**] that there is a corresponding finite obstruction set $\mathcal{O}_n$. For each n, $\mathcal{O}_n$ is the set of graphs that are not n-apex even though every proper minor is. We say that the elements of $\mathcal{O}_n$ are **minor minimal not n-apex** or MMNnA.

Determining this obstruction set is likely to be quite difficult. It's known that, already, $|\mathcal{O}_1| > 150$ [**P**], and we expect that $|\mathcal{O}_n|$ grows quickly in size with n. In the current paper we report on computer searches that show $|\mathcal{O}_2| \geq 82$, $|\mathcal{O}_3| \geq 601$, $|\mathcal{O}_4| \geq 520$, and $|\mathcal{O}_5| \geq 608$.



Figure 1. ∇Y and Y∇ moves.

We obtain these bounds on the size of $\mathcal{O}_n$ using the Y∇Y families of K_{n+5}, the complete graph on $n+5$ vertices, and $K_{3^2,1^n}$, the complete multi-partite graph

2000 *Mathematics Subject Classification.* 05C10.

©0000 (copyright holder)

with two parts of three vertices and a further n parts that each have a single vertex. As in Figure 1, we perform a ∇Y or **triangle-Y move** on a graph by deleting the edges of a triangle abc and then adding a vertex v and edges av, bv, and cv. The reverse operation is a Y∇ or **Y-triangle move**. The set of graphs obtained from a graph G through a sequence of zero or more ∇Y or Y∇ moves is G**'s (**Y∇Y**) family**.

In this paper we provide evidence in support of the following conjecture.

CONJECTURE 1. *For each $n > 0$, the K_{n+5} and $K_{3^2,1^n}$ families are in $\mathcal{O}_n$.*

When $n = 0$, there are 49 graphs in the K_5 family and 10 in $K_{3,3}$'s. However, $\mathcal{O}_0 = \{K_5, K_{3,3}\}$ as proved by Weber [**W**] and the analogue of the conjecture does not hold in this case. On the other hand, a proof for $n \leq 2$ follows easily from earlier work.

PROPOSITION 2. *The conjecture holds for $n \leq 2$.*

PROOF. For $n = 1$, the K_6 and $K_{3,3,1}$ families coincide. This collection of seven graphs is known as the Petersen family since it also includes the Petersen graph. Barsotti and Mattman [**BM**] observed that this family is in $\mathcal{O}_1$ on the way to proving that the K_7 family is in $\mathcal{O}_2$. In fact, they showed that the K_7 family is exactly the set of graphs in $\mathcal{O}_2$ of size 21 or less.

It remains to show that $K_{3^2,1^2}$ is in $\mathcal{O}_2$. In [**GMN**], Goldberg, Mattman, and Naimi show that the 58 graphs in this family are all minor minimal intrinsically knotted or MMIK. This implies they are not 2-apex, see [**BBFFHL, OT**]. Since K_7 and its 13 descendants are also MMIK [**KS**], this means that none of these 14 graphs can be a minor of a $K_{3^2,1^2}$ family graph.

So far, we have established that each of the size 22 graphs in the $K_{3^2,1^2}$ family is not 2-apex. If one of these were not in $\mathcal{O}_2$, then it would have to have a 2-apex minor of size at most 21. By [**BM**], this minor would lie in the K_7 family. We've already argued that 14 of the 20 graphs in that family cannot play this role. We omit the straightforward verification that none of the remaining six graphs in the K_7 family is a minor of a member of the $K_{3^2,1^2}$ family. □

We verify the conjecture for $n = 3$ with the aid of a computer, as described in Section 2 below. For $n = 4$, we checked that the 163 graphs in the K_9 family and 357 of the graphs in the $K_{3^2,1^4}$ family (all those within four ∇Y or Y∇ moves of $K_{3^2,1^4}$) are in $\mathcal{O}_4$. For $n = 5$, the computer assures us that 608 of the 1681 graphs in K_{10}'s family are in $\mathcal{O}_5$. This gives the bounds for $\mathcal{O}_4$ and $\mathcal{O}_5$ mentioned earlier. Due to time constraints, we did not attempt to check more graphs in the K_{10} family nor in the $K_{3^2,1^4}$ family of more than 13 thousand graphs.

It is straightforward to see that K_{n+5} and $K_{3^2,1^n}$ themselves are in $\mathcal{O}_n$. We also verify this for the three immediate descendants of each of these two graphs in the following theorem, which we prove in Section 1. Recall that when a sequence of ∇Y moves take us from G_1 to G_2, we say G_2 is a **descendant** of G_1.

THEOREM 3. *Let $n > 0$. The graphs K_{n+5} and $K_{3^2,1^n}$ and the three immediate descendants of each are elements of $\mathcal{O}_n$*

It would be tempting to argue that ∇Y and Y∇ moves preserve the MMNnA property that characterizes $\mathcal{O}_n$. Unfortunately, this is not true in general. For example, the disjoint unions $G_3 = \sqcup_{i=1}^{n+1} K_{3,3}$ and $G_5 = \sqcup_{i=1}^{n+1} K_5$ are both MMNnA.

However, a Y∇ move on $K_{3,3}$ results in a planar graph and, similarly, a Y∇ move on G_3 gives a graph that is not in $\mathcal{O}_n$. On the other hand, a ∇Y move on K_5 results in a graph that is nonplanar, but with proper $K_{3,3}$ minor. So, the child of G_5 cannot join it in $\mathcal{O}_n$.

On the other hand, it's easy to see that Y∇ moves preserve planarity (or, equivalently, ∇Y preserves nonplanarity). If G is planar, and a Y∇ move on the degree 3 vertex v results in G' (We call this a Y∇ **move at** v.) then, fixing a planar embedding of G, we can effect the Y∇ move in a tubular neighborhood of v and its three edges to give a planar embedding of G'.

Using this idea we can show that a Y∇ move at v preserves the n-apex property provided v is not an apex. For 1-apex graphs, this observation is due to Barsotti in unpublished work. For $W \subset V(G)$, let $G - W$ denote the induced subgraph on $V(G) \setminus W$.

LEMMA 4. *Suppose G is n-apex with subset U of $V(G)$ such that $|U| \leq n$ and $G - U$ is planar. If G' is obtained from G by a* Y∇ *move at the vertex $v \notin U$ then G' is also n-apex.*

PROOF. The key observation is that the Y∇ move preserves planarity so that $G' - U$ is again planar and G' is $|U|$-apex. The details depend on how many of v's neighbors, a, b, and c, are in U.

If none are, the argument is the same as above: we can effect the Y∇ move in a tubular neighborhood of v and its edges. If exactly one is, say $a \in U$, then the Y∇ move replaces the edges vb and vc in $G - U$ with the edge bc in $G' - U$ so that planarity is preserved. Similarly, if two or all three neighbors are in U, the Y∇ move will leave $G' - U$ planar. □

Although Y∇ and ∇Y moves do not preserve MMNnA in general, our conjecture asserts that they do in the K_{n+5} and $K_{3^2,1^n}$ family.

We describe results for $\mathcal{O}_2$ and $\mathcal{O}_3$ in Section 2. This includes classifications of all the graphs in $\mathcal{O}_2$ through order 9 and size 23. We also give a proof "by hand" that the 32 graphs in the K_8 family are in $\mathcal{O}_3$. This, with the computer verification for the $K_{3^2,1^3}$ family, shows that our conjecture holds for $n = 3$. But first we prove Theorem 3 in the next section.

1. Proof of Theorem 3

In this section we prove that K_{n+5}, $K_{3^2,1^n}$, and their three immediate descendants are in $\mathcal{O}_n$. In the introduction, we gave the proof for $n \leq 2$, so we will assume $n \geq 3$. Then, the three descendants of $K_{3^2,1^n}$ are its **children**, the graphs obtained by a single ∇Y move. The graph K_{n+5} has a single child, C_1, which in turn has two children. This child and pair of grandchildren are the three immediate descendants of K_{n+5} referred to in the statement of Theorem 3.

We begin with a lemma that will be useful for one of the children of $K_{3^2,1^n}$ and which gives the flavor of our approach.

LEMMA 5. *Let D_3'' denote the complement of the $(12, 17)$ graph of Figure 2 and D_3' the induced subgraph on $V(D_3'') \setminus \{v_{10}, v_{11}\}$. Then $D_3' \in \mathcal{O}_3$ and $D_3'' \in \mathcal{O}_5$.*

Since D_3' is a child of $K_{3^2,1^3}$ and D_3'' is a child of $K_{3^2,1^5}$, the lemma is consistent with our conjecture.



FIGURE 2. The complement of the graph D_3''.

PROOF. We argue D_3' is not 3-apex by showing how deleting any three vertices results in a nonplanar graph. Deleting v_0 yields $D_3' - v_0 = K_{3^3}$, which is not 2-apex. So we can assume that v_0 is not among the three deleted vertices. On the other hand, if none of $\{v_7, v_8, v_9\}$ are deleted, then a $K_{3,3}$ subgraph remains after deleting three vertices from $\{v_1, v_2, v_3, v_4, v_5, v_6\}$. Yet, if $\{v_7, v_8, v_9\}$ are the three deleted vertices, again a $K_{3,3}$ subgraph remains. So, we are left with the cases where exactly one or two vertices in $\{v_7, v_8, v_9\}$ is deleted.

Suppose then that v_9 is not deleted. Since we are assuming v_0 is also not deleted, we can contract v_0v_9. Then, deleting v_7 leaves the graph $K_{3^2,1^2}$, which is not 2-apex. This completes the argument that D_3' is not 3-apex.

To show that D_3' is minor minimal not 3-apex, we argue that for any $e \in E(D_3')$, both deletion, $D_3' - e$, and contraction, D_3'/e, of that edge gives a 3-apex graph. Up to symmetry there are three types of edges: v_0v_7, v_1v_4, v_1v_7. In each case, we list three vertices in an apex set.

Edge e	Apex Sets
v_0v_7	$D_3' - e$: v_1, v_4, v_8; D_3'/e: $\overline{v_0}, v_1, v_4$
v_1v_4	$D_3' - e$: v_2, v_3, v_7; D_3'/e: v_2, v_3, v_7
v_1v_7	$D_3' - e$: v_4, v_5, v_8; D_3'/e: v_2, v_3, v_8

For example, when $e = v_0v_7$, $D_3' - e$ becomes planar on deletion of vertices v_1, v_4, and v_8. We use $\overline{v_0}$ to denote the vertex of D_3'/v_0v_7 that comes from identifying v_0 and v_7. This complete the proof that $D_3' \in \mathcal{O}_3$.

Our argument for D_3'' is similar and we make use of what was proved for the subgraph D_3'. We show that deleting five vertices from D_3'' leaves a nonplanar graph. If both v_{10} and v_{11} are among the five, this is true as deleting those two gives D_3', which is not 3-apex as we just showed. This leaves two cases: either exactly one or else neither of v_{10} and v_{11} is deleted.

If neither is deleted, note that those two together with one vertex each from $\{v_1, v_2, v_3\}$, $\{v_4, v_5, v_6\}$, and $\{v_7, v_8, v_9\}$ induce a K_5 subgraph. So, we can assume that one of those triples is entirely deleted. For example, say the five deleted vertices include $\{v_1, v_2, v_3\}$. Then at most two of $\{v_7, v_8, v_9\}$ are deleted. Say v_7 is not. Contracting v_0v_7 gives a subgraph of $K_{3^2,1^2}$ which is not 2-apex. Suppose instead that $\{v_7, v_8, v_9\}$ is included among the five deleted vertices. If v_0 is also deleted, what remains is $K_{3^2,1^2}$, which not only fails to be 1-apex (which is all that

is required for our argument), but is in fact not 2-apex. If v_0 is not among the deleted vertices, then, up to symmetry, there are only two ways to choose the last two vertices to delete: either $\{v_1, v_2\}$ or $\{v_1, v_4\}$. In either case, a nonplanar graph results.

To complete the argument that D_3'' is not 5-apex, suppose that exactly one of v_{10} and v_{11} is deleted, say v_{11}. If we, in addition, delete v_0, what remains is a $K_{3^3,1}$ graph. There are then three vertices left to delete, and if any pair are in the same part of $K_{3^3,1}$, after deleting that pair, we have the not 1-apex graph $K_{3^2,1}$ as a subgraph and a nonplanar graph on deleting the fifth and final vertex. So, we can assume that the remaining three deleted vertices are taken one each from different parts of $K_{3^3,1}$. For example, we could delete v_1, v_4, and v_7. This leaves a nonplanar graph.

This means we can assume v_0 is not among the five deleted vertices, still under the assumption that v_{11}, but not v_{10}, is deleted. If v_1 and v_2 are deleted, the remaining graph has a $K_{3^2,1^2}$ subgraph and is not 2-apex. Using the symmetry of the graph, this means we can assume v_7 and v_8 are deleted instead. However, this again results in a graph that is not 2-apex by virtue of having a $K_{3^2,1^2}$ subgraph. This completes the argument that D_3'' is not 5–apex.

We next demonstrate that, for any edge e, both $D_3'' - e$ and D_3''/e are 5-apex. If e is not incident on v_{10} or v_{11}, we may delete those two vertices and recognize e as an edge in D_3'. As we have already found three vertex apex sets for $D_3' - e$ and D_3'/e, taking the union with $\{v_{10}, v_{11}\}$ gives a five vertex apex set for $D_3'' - e$ or D_3''/e.

So, we may assume that e is incident on v_{10}, v_{11} or both. Up to symmetry, there are three such edges, and, in each case, we provide a five vertex apex set.

Edge e	Apex Sets
v_1v_{10}	$D_3'' - e$: v_4, v_5, v_6, v_7, v_8; D_3''/e: v_2, v_3, v_4, v_7, v_8
v_7v_{10}	$D_3'' - e$: $v_1, v_4, v_8, v_9, v_{11}$; D_3''/e: v_1, v_2, v_3, v_4, v_8
$v_{10}v_{11}$	$D_3'' - e$: v_1, v_4, v_7, v_8, v_9; D_3''/e: v_1, v_2, v_3, v_4, v_7

This completes the proof that $D_3'' \in \mathcal{O}_5$. □

PROOF. (of Theorem 3) We begin with K_{n+5}, which is not n-apex since deletion of any n vertices gives the nonplanar graph K_5. On the other hand, given $e \in E(K_{n+5})$, both deletion and contraction of e result in an n-apex graph as can be seen by deleting any n vertices not on the edge e. Therefore, every proper minor of K_{n+5} is n-apex and $K_{n+5} \in \mathcal{O}_n$.

Let C_1 denote the child of K_{n+5}. This graph has a unique vertex, v_0, of degree three with neighbors v_1, v_2, and v_3. Denote the remaining vertices $v_4, v_5, v_6, \ldots, v_{n+5}$. Generally, deleting n vertices results in a graph with $K_{3,3}$ minor. The exception is when exactly one of v_1, v_2, and v_3 is deleted and v_0 is not. In this case, the result has a K_5 minor. Thus, C_1 is not n-apex.

Up to symmetry, there are three types of edges in C_1, represented by v_0v_1, v_1v_4, and v_4v_5. For each choice of edge, e, we argue that both deletion, $C_1 - e$, and contraction, C_1/e, gives an n-apex graph.

For C_1/e, we observe that, no matter which edge is contracted, by appropriate deletion of n vertices, we can ensure that v_0 (or the vertex formed by identifying v_0

and v_1 in case $e = v_0v_1$) is an isolated vertex in the resulting graph on 5 vertices, which is, therefore, nonplanar. If $e = v_0v_1$ or v_1v_4, $C_1 - e$ is planar after deleting all vertices but v_0 through v_6, while $C_1 - v_4v_5$ becomes planar after deletion of at least one of v_1, v_2, v_3 and none of v_0, v_4, v_5.

Therefore $C_1 \in \mathcal{O}_n$ and we turn to its children. To form a triangle in C_1, we use at most one vertex from v_1, v_2, and v_3. This means a triangle cannot include v_0 and, up to symmetry, there are two types of triangles: $v_1v_4v_5$, resulting in the child C_2, or $v_4v_5v_6$, which gives a child we'll call C_3. For these new graphs, we'll use the same vertex labels as in C_1 with the addition of a new degree three vertex w_0.

Let's see why $C_2 \in \mathcal{O}_n$. First, notice that there is a graph automorphism by interchanging v_0 and w_0 and the pair $\{v_2, v_3\}$ with $\{v_4, v_5\}$. Deleting $n-2$ of the n vertices in $\{v_6, v_7, \dots v_{n+5}\}$ results in a graph in the K_7 family, that is not 2-apex by Proposition 2. The only way C_2 could remain planar after deletion of n vertices is if at least three of those vertices remain. However the remaining vertices from that set, together with up to one each of $\{v_2, v_3\}$ and $\{v_4, v_5\}$, form a clique. A clique on five or more vertices contains K_5 and is nonplanar. Thus, we surely have a nonplanar subgraph by deletion of n vertices except, possibly, in two cases (up to symmetry): 1) $\{v_6, v_7, v_8, v_9\}$ remain after deleting the n vertices, or 2) $\{v_6, v_7, v_8\}$ and vertices from at most one of $\{v_2, v_3\}$ and $\{v_4, v_5\}$, say the first, remain.

In the first case, we see that $\{v_2, v_3, v_4, v_5, v_{10}, v_{11}, \dots v_{n+5}\}$ are the n deleted vertices and what remains is the graph induced on v_0, v_1, v_6, v_7, v_8, v_9, and w_0, which is nonplanar.

In the second case, we can assume v_2 remains as otherwise we must have deleted the $n+1$ vertices $\{v_2, v_3, v_4, v_5, v_9, v_{10}, \dots, v_{n+5}\}$. Since v_2 remains, we see that $\{v_4, v_5, v_9, v_{10}, \dots, v_{n+5}\}$ are $n-1$ of the deleted vertices. The graph H that remains after deleting these is the induced graph on the eight vertices v_0, v_1, v_2, v_3, v_6, v_7, v_8, and w_0. It is straightforward to verify that H is not 1-apex. So, no matter which vertex is chosen as the nth for deletion, the result will be nonplanar in this case. This completes the argument that C_2 is not n-apex.

Next we show that every proper minor is n-apex by establishing this for $C_2 - e$ and C_2/e whatever edge e is chosen. Up to symmetry, there are six choices for e: v_0v_1, v_0v_2, v_1v_6, v_2v_4, v_2v_6, and v_6v_7. Recall that deleting $n-2$ vertices from $\{v_6, v_7, \dots, v_{n+5}\}$ converts C_2 to a graph C_2' in the K_7 family. Each choice of e corresponds to an edge in C_2' and we've already established, in Proposition 2, that $C_2' - e$ and C_2'/e are 2-apex. This means the corresponding minors of C_2 are n-apex. This completes the argument that $C_2 \in \mathcal{O}_n$.

Let's see why C_3 is not n-apex. Deleting $n-2$ vertices in $V_{7+} = \{v_7, v_8, \dots, v_{n+5}\}$ results in a graph in the K_7 family that is not 2-apex by Proposition 2. So, to achieve a planar graph after deleting n vertices would require at least two vertices in V_{7+} to remain. On the other hand the vertices in V_{7+} induce a clique so, to avoid the nonplanar K_5, at most four can remain. Up to symmetry, this gives three cases: 1) $\{v_7, v_8, v_9, v_{10}\}$, 2) $\{v_7, v_8, v_9\}$, or 3) $\{v_7, v_8\}$ are the elements that remain after deleting n–vertices. To complete the argument that C_3 is not n-apex, we'll show that none of these lead to a planar graph.

Any vertex from $\{v_1, v_2, v_3\}$ or $\{v_4, v_5, v_6\}$ will also induce a clique with vertices from V_{7+}. In the first case, if we are to avoid a nonplanar graph, we must delete the $n+1$ vertices $\{v_1, v_2, \dots v_6, v_{11}, v_{12}, \dots, v_{n+5}\}$, so we cannot use this case to show C_3

is n-apex. In the second case, we could have vertices from at most one of $\{v_1, v_2, v_3\}$ and $\{v_4, v_5, v_6\}$, say the first, in addition to $\{v_7, v_8, v_9\}$. This means we have $\{v_4, v_5, v_6, v_{10}, v_{11}, \ldots, v_{n+5}\}$ as $n-1$ of the deleted vertices. It's straightforward to verify that the graph induced on the other vertices, $v_0, v_1, v_2, v_3, v_7, v_8, v_9, w_0$ is not 1-apex. For the final case, $\{v_9, v_{10}, \ldots, v_{n+5}\}$ are $n-3$ of the deleted vertices. The graph induced on the other vertices, $v_0, v_1, \ldots v_8, w_0$ is not 3-apex. Thus, in any case, we deduce that C_3 is not n-apex.

Up to symmetry, there are four types of edges in C_3: v_0v_1, v_1v_4, v_1v_7, and v_7v_8. By deleting $n-2$ vertices in $\{v_7, v_8, \ldots, v_{n+5}\}$ we arrive at the graph C_3' in the K_7 family. Aside from v_7v_8, each choice of edge e is represented in C_3'. In Proposition 2 we argued that $C_3' - e$ and C_3'/e are 2-apex, which means $C_3 - e$ and C_3/e are n-apex. When $e = v_7v_8$, we can think of deleting the $n-3$ vertices $\{v_9, v_{10}, \ldots, v_{n+5}\}$ to give the graph H induced on the vertices $v_0, v_1, \ldots v_8, w_0$, a graph in the K_8 family. Now, deleting vertices v_1, v_2, v_4 shows that $H - v_7v_8$ is 3-apex while deleting vertices v_1, v_2, v_3 demonstrates that H/v_7v_8 is also 3-apex. It follows that deleting or contracting v_7v_8 in C_3 results in an n-apex graph. Since any $C_3 - e$ or C_3/e is n-apex, it follows that any proper minor is n-apex. This completes the argument that $C_3 \in \mathcal{O}_n$.

The graph $K_{3^2,1^n}$ is not n-apex as deleting any n vertices leaves a graph with $K_{3,3}$ minor. Let v_1, v_2, v_3 and v_4, v_5, v_6 denote the vertices in the two parts with three vertices and $v_7, v_8, \ldots v_{n+6}$ the vertices in the remaining n parts. Up to symmetry, there are three types of edges e: v_1v_4, v_1v_7, and v_7v_8. Removing n vertices from $K_{3^2,1^n}/e$ gives a graph on five vertices which is planar as long as it is not K_5. For example, ensuring that v_5 and v_6 are among the five vertices that remain shows that $K_{3^2,1^n}/e$ is n-apex. On the other hand, deleting n vertices from $K_{3^2,1^n} - e$ gives a graph on six vertices. For each choice of e, we list six vertices that induce a planar subgraph: for $e = v_1v_4$, use $v_1, v_2, \ldots, v_6$; if $e = v_1v_7$, take $v_1, v_2, v_3, v_4, v_5, v_7$; and when $e = v_7v_8$, then $v_1, v_2, v_4, v_5, v_7, v_8$ will work. This shows that any proper minor of $K_{3^2,1^n}$ is n-apex and completes the proof that it is in $\mathcal{O}_n$.

Up to symmetry, there are three types of triangles in $K_{3^2,1^n}$. We'll call the three children D_1, D_2, and D_3, corresponding to ∇Y moves on on $v_1v_4v_7$, $v_1v_7v_8$, and $v_7v_8v_9$ respectively. We'll argue each child is in $\mathcal{O}_n$. In D_i, we preserve the vertex labels $v_1, v_2, \ldots, v_{n+6}$ from $K_{3^2,1^n}$ and call the new vertex of degree three v_0.

In D_1, deleting $n-2$ vertices from $V_{8+} = \{v_8, v_9, \ldots, v_{n+6}\}$ gives a graph, D_1', in the $K_{3^1,1^2}$ family that is not 2-apex by Proposition 2. So, at least two vertices from V_{8+} must remain. On the other hand, vertices in V_{8+} induce a clique so at most four can remain and we again have three cases: 1) $\{v_8, v_9, v_{10}, v_{11}\}$, 2) $\{v_8, v_9, v_{10}\}$, and 3) $\{v_8, v_9\}$ are the vertices of V_{8+} that remain.

In addition, up to one vertex each from $\{v_2, v_3\}$ and $\{v_5, v_6\}$ can be added to those remaining from V_{8+} to make a clique. So, in the first case, we can assume that $n-1$ vertices were deleted leaving the graph induced on $v_0, v_1, v_4, v_7, v_8, v_9, v_{10}, v_{11}$, which is not 1-apex. In the second case, let's say that vertices from $\{v_2, v_3\}$ may remain. Then, after deleting $n-2$ vertices, we have the graph induced on $v_0, v_1, v_2, v_3, v_4, v_7, v_8, v_9, v_{10}$, which is not 2-apex. Finally, in the third case, after deleting $n-3$ vertices, we have the graph induced on $v_0, v_1, \ldots, v_9$, which is not 3-apex. This shows that D_1 is not n-apex.

Up to symmetry there are nine types of edges in D_1: v_0v_1, v_0v_7, v_1v_5, v_1v_8, v_2v_5, v_2v_7, v_2v_8, v_7v_8, and v_8v_9. Except for the last one, v_8v_9, each edge e is represented in D_1', a graph in the $K_{3^2,1^2}$ family obtained by deleting $n-2$ vertices from V_{8+}. In Proposition 2, we argued that $D_1'-e$ and D_1'/e are 2-apex and it follows that D_1-e and D_1/e are n-apex when $e \neq v_8v_9$. If we delete $n-3$ vertices in V_{8+}, leaving $\{v_8, v_9\}$, we obtain the graph H induced on $v_0, v_1, \ldots, v_9$, which includes the edge v_7v_8. Deleting v_1, v_4, v_7 shows that $H - v_8v_9$ is 3-apex and $D_1 - v_8v_9$ is n-apex. For H/v_8v_9, deletion of v_1, v_2, v_3 shows that it it 3-apex and D_1/v_8v_9 is n-apex. We've shown that for any edge e, $D_1 - e$ and D_1/e are both n-apex. It follows that any proper minor is n-apex, which completes the argument that $D_1 \in \mathcal{O}_n$.

The verification for D_2 is similar and we'll omit some details. Deleting the $n-2$ vertices in V_{9+} results in the graph D_2' in the $K_{3^2,1^2}$ family, which is not 2-apex. This leaves the cases where between four and one vertex of V_{9+} remain. Since one vertex each from $\{v_2, v_3\}$, $\{v_4, v_5, v_6\}$, and $\{v_7, v_8\}$ can join with vertices of V_{9+} to form a clique, some or all of these must be deleted to avoid K_5, a clique on five vertices, surviving. For example, in the case that four vertices of V_{9+} remain, then all seven vertices $v_2, v_3 \ldots v_8$ must be deleted in addition to the $n-6$ taken from V_{9+}, which is a total of $n+1$ vertices. In the case that only one vertex of V_{9+} remains, we have removed $n-3$ vertices from V_{9+} and argue that the graph that remains, on ten vertices, is not 3-apex. The remaining cases can be approached in the same way; remove either $n-4$ or $n-5$ vertices from V_{9+} and argue that the remaining graphs are not 4-apex nor 5-apex. This shows D_2 is not n-apex.

There are several edge types in D_2, but with the exception of those incident on vertices in V_{9+}, all are represented in D_2'. For edges e incident on one or two vertices of V_{9+}, we argue directly that the graphs that remain after deleting $n-3$ or $n-4$ vertices of V_{9+} and either contracting or deleting e are 3-apex or 4-apex, respectively. This implies that the corresponding graph $D_2 - e$ or D_2/e is n-apex, as required.

Finally, we turn to D_3. As with earlier cases, we will leverage a similar graph D_3', this time in the $K_{3^2,1^3}$ family, In Lemma 5 we proved $D_3' \in \mathcal{O}_3$. Deleting the $n-3$ vertices in V_{10+} gives D_3', so to prove D_3 is not n-apex it remains to take the cases where between one and four vertices of V_{10+} remain.

Taking a vertex each from $\{v_1, v_2, v_3\}$, $\{v_4, v_5, v_6\}$, and $\{v_7, v_8, v_9\}$ gives a clique with any vertices that remain from V_{10+}. In case four vertices from V_{10+} remain, this means deletion of a further nine vertices in addition to the $n-7$ from V_{10+} to avoid a K_5 minor and so cannot be used to show D_3 is n-apex. Suppose the three vertices $\{v_{10}, v_{11}, v_{12}\}$ remain from V_{10+}. To avoid a K_5 minor, we must delete two of the triples and there are a couple of ways to do this. If $\{v_1, v_2, \ldots, v_6\}$ are deleted, then n vertices are gone and we observe that the resulting graph induced on $v_0, v_7, v_8, \ldots, v_{12}$ is nonplanar. On the other hand, if $\{v_4, v_5, \ldots, v_9\}$ are deleted, the result, induced on $v_0, v_1, v_2, v_3, v_{10}, v_{11}, v_{12}$ is again nonplanar. So, we can assume at most two vertices from V_{10+} remain. If two, we've deleted $n-5$ vertices to leave the graph $D_3'' \in \mathcal{O}_5$ (see Lemma 5) and if only one, what's left is a graph in the $K_{3^2,1^6}$ family that can be shown to be not 6-apex. This completes the argument that D_3 is not n-apex.

Finally, any $e \in E(D_3)$ is realized as an edge of D_3''. In Lemma 5, we prove that $D_3'' - e$ and D_3''/e are 5-apex and it follows that D_3 and D_3'' are n-apex, completing the proof that $D_3 \in \mathcal{O}_n$. □

2. Results for $\mathcal{O}_2$ and $\mathcal{O}_3$

In this section we prove Conjecture 1 when $n = 3$. However, we begin with classifications of the graphs in $\mathcal{O}_2$ through order nine and size 23. Mattman [**M**] established that the elements of $\mathcal{O}_2$ have size 21 at least and Barsotti and Mattman [**BM**] showed that the K_7 family is precisely the set of graphs of size 21. The 58 $K_{3^2,1^2}$ graphs are of size 22 in $\mathcal{O}_2$ and, based on a computer search, we have found exactly two other graphs with 22 edges. Both are 4–regular on 11 vertices and we list their edges below.

$$\{\{1,6\},\{1,9\},\{1,10\},\{1,11\},\{2,6\},\{2,9\},\{2,10\},\{2,11\},$$
$$\{3,7\},\{3,8\},\{3,9\},\{3,10\},\{4,7\},\{4,8\},\{4,9\},$$
$$\{4,11\},\{5,7\},\{5,8\},\{5,10\},\{5,11\},\{6,7\},\{6,8\}\}$$

$$\{\{1,5\},\{1,6\},\{1,7\},\{1,8\},\{2,6\},\{2,7\},\{2,8\},\{2,9\},$$
$$\{3,7\},\{3,8\},\{3,9\},\{3,10\},\{4,8\},\{4,9\},\{4,10\},$$
$$\{4,11\},\{5,9\},\{5,10\},\{5,11\},\{6,10\},\{6,11\},\{7,11\}\}$$

We were surprised to discover that there are **no** graphs in $\mathcal{O}_2$ of size 23. This is similar to $\mathcal{O}_1$ where the Petersen family graphs of size 15 are the only obstructions with fewer than 18 edges (see [**BM**]). On the other hand, we do have examples of graphs in $\mathcal{O}_2$ for each size between 24 and 30, inclusive.

In terms of order, we found a total of 12 graphs in $\mathcal{O}_2$ with nine or fewer vertices. There are five each in the K_7 and $K_{3^2,1^2}$ families and two of size 26 and 27 whose edge lists we give below.

$$\{\{1,4\},\{1,5\},\{1,7\},\{1,8\},\{1,9\},\{2,5\},\{2,6\},\{2,7\},\{2,8\},$$
$$\{2,9\},\{3,5\},\{3,6\},\{3,7\},\{3,8\},\{3,9\},\{4,6\},\{4,7\},\{4,8\},$$
$$\{4,9\},\{5,6\},\{5,8\},\{5,9\},\{6,8\},\{6,9\},\{7,8\},\{7,9\}\}$$

$$\{\{1,4\},\{1,5\},\{1,6\},\{1,7\},\{1,8\},\{1,9\},\{2,4\},\{2,5\},\{2,6\},$$
$$\{2,7\},\{2,8\},\{2,9\},\{3,4\},\{3,5\},\{3,6\},\{3,7\},\{3,8\},\{3,9\},$$
$$\{4,6\},\{4,7\},\{4,8\},\{5,7\},\{5,8\},\{5,9\},\{6,8\},\{6,9\},\{7,9\}\}$$

In total, this gives 82 graphs in $\mathcal{O}_2$: 20 in the K_7 family, 58 in $K_{3^2,1^2}$'s, and four more with edges listed above.

Next, we show that Conjecture 1 holds for $n = 3$. For this we rely on a computer verification that the 569 graphs in the $K_{3^2,1^3}$ family are MMN3A. Then the following theorem completes the proof. Together with the $K_{3^2,1^3}$ family, the 32 graphs in K_8's give the bound $\mathcal{O}_3 \geq 601$ mentioned in the introduction.

THEOREM 6. *The K_8 family is in $\mathcal{O}_3$.*

PROOF. The K_8 family consists of 32 graphs, illustrated schematically in Figure 3. In the appendix we give edge lists for these graphs. Graphs at the same height are of the same order and edges join graphs that are related by a ∇Y move. All graphs in this family have 28 edges. The numbering simply reflects the order in which these graphs were generated by our computer program and we refer to the graphs in the family as Cousin 1, K_8, through Cousin 32, a graph of order 16.



FIGURE 3. The K_8 family.

Theorem 3 shows that Cousin 1, K_8, and its child, Cousin 2, and grandchildren, Cousins 3 and 4, are all MMN3A. We apply Lemma 4 to see that the remaining descendants of K_8 are not 3-apex.

For example, let C_5 denote Cousin 5, which is a child of Cousin 3, C_3. In other words a Y∇ move at some vertex v in C_5 gives C_3. If C_5 were 3-apex and v not an apex, then Lemma 4 would imply C_3 is 3-apex, a contradiction. So, to show that C_5 is not 3-apex, it is enough to argue that v cannot be an apex. For this, it suffices to show that $C_5 - v$ is not 2-apex. Indeed, C_3 can be obtained by a Y∇ move at vertex $v = 9$. Deleting edges $\{4, 6\}$, $\{4, 8\}$, and $\{6, 8\}$ from $C_5 - 9$ results in a graph (Cousin 4 by the numbering in [**GMN**]) in the $K_{3^2,1^2}$ family. This shows that $C_5 - 9$ is not 2-apex, as required.

In the appendix, we list, for each cousin C_n ($5 \leq n \leq 32$) in the K_8 family, a vertex v such that a Y∇ move at v gives another graph in the family, and edges of C_n such that deleting or contracting those edges from $C_n - v$ gives a graph in the $K_{3^2,1^2}$ family. In fact, we can take $v = 9$ in all cases. This shows no graphs in the family are 3-apex.

Further, for $5 \leq n \leq 32$, we show that each proper minor of C_n is 3-apex. For each edge e (up to symmetry), we show that both $C_n - e$ and C_n/e (edge deletion

and contraction, respectively) are 3-apex by listing three vertices whose deletion gives a planar subgraph. This completes the proof. □

Appendix A. Edge lists of K_8 family graphs

In Tables 1 and 2 we list the edges of the 32 graphs in the K_8 family. We number the graphs as Cousin 1 (K_8) through Cousin 32 (a graph of order 16).

Appendix B. K_8 family graphs are not 3-apex

For each cousin C_n ($5 \leq n \leq 32$) in the K_8 family, a Y∇ move at vertex 9 results in another graph in that family. We can show C_n is not 3-apex by arguing that $C_n - 9$ is not 2-apex. For this we list, in Table 3, edges that are to be deleted or contracted in $C_n - 9$ to produce a minor which is a cousin in the $K_{3^2,1^2}$ family. Cousins in that family are numbered as in [**GMN**]

For example, deleting edges $\{\{4,6\},\{4,8\},\{6,8\}\}$ of $C_5 - 9$ results in cousin 4 of the $K_{3^2,1^2}$ family. In the case of $C_8 - 9$, first delete edges $\{4,6\}$ and $\{4,8\}$ and then contract edge $\{8,11\}$ to realize cousin 2 of the $K_{3^2,1^2}$ family.

Appendix C. Proper minors are 3-apex

For each cousin C_n ($5 \leq n \leq 32$) in the K_8 family, each proper minor of C_n is 3-apex. We show this in Tables 4–10 by considering each edge e of C_n (distinct up to symmetry) and for the graphs $C_n - e$ and C_n/e (edge deletion and edge contraction, respectively) listing three vertices whose deletion gives a planar subgraph.

In Tables 4–10 we use the convention that the vertex that is the result of an edge contraction adopts the name of the smaller of the two vertex names on the edge that was contracted. For example if we were to contract the edge $\{1,8\}$, the new resulting vertex would be referred to as vertex 1.

References

[BM] J. Barsotti and T.W. Mattman, *Graphs on 21 edges that are not 2-apex.* (To appear in Involve). arXiv:1506.06789

[BBFFHL] P. Blain, G. Bowlin, T. Fleming, J. Foisy, J. Hendricks, and J. LaCombe. *Some Results on Intrinsically Knotted Graphs.* J. Knot Theory Ramifications, **16** (2007), 749–760.

[GMN] N. Goldberg, T.W. Mattman, and R. Naimi. Many, many more intrinsically knotted graphs, *Algebr. Geom. Topol.* **14** (2014), 1801-1823. arXiv:1109.1632

[KS] T. Kohara and S. Suzuki. *Some remarks on knots and links in spatial graphs.* Knots 90, Osaka, 1990, de Gruyter (1992) 435–445.

[K] K. Kuratowski. Sur le problème des courbes gauches en topologie. *Fund. Math.* **15** (1930) 271–283.

[M] T. W. Mattman. *Graphs of 20 edges are 2-apex, hence unknotted.* Alg. Geom. Top., **11** (2011) 691–718. arxiv.org/0910.1575

[OT] M. Ozawa and Y. Tsutsumi. *Primitive Spatial Graphs and Graph Minors.* Rev. Mat. Complut., **20** (2007), 391–406.

[P] M. Pierce, *Searching for and classifying the finite set of minor-minimal non-apex graphs.* CSU, Chico Senior Thesis, (2014). Available at `http://www.csuchico.edu/~tmattman`.

[RS] N. Robertson, P. Seymour. *Graph minors. XX. Wagner's conjecture.* J. Combin. Theory Ser. B, **92**, (2004), 325–357.

[W] K. Wagner. Über eine Eigenschaft der ebenen Komplexe. *Math. Ann.* **114** (1937), 570–590.

n	Edges of C_n
1	{{1, 2}, {1, 3}, {1, 4}, {1, 5}, {1, 6}, {1, 7}, {1, 8}, {2, 3}, {2, 4}, {2, 5}, {2, 6}, {2, 7}, {2, 8}, {3, 4}, {3, 5}, {3, 6}, {3, 7}, {3, 8}, {4, 5}, {4, 6}, {4, 7}, {4, 8}, {5, 6}, {5, 7}, {5, 8}, {6, 7}, {6, 8}, {7, 8}}
2	{{1, 4}, {1, 5}, {1, 6}, {1, 7}, {1, 8}, {1, 9}, {2, 4}, {2, 5}, {2, 6}, {2, 7}, {2, 8}, {2, 9}, {3, 4}, {3, 5}, {3, 6}, {3, 7}, {3, 8}, {3, 9}, {4, 5}, {4, 6}, {4, 7}, {4, 8}, {5, 6}, {5, 7}, {5, 8}, {6, 7}, {6, 8}, {7, 8}}
3	{{1, 6}, {1, 7}, {1, 8}, {1, 9}, {1, 10}, {2, 4}, {2, 5}, {2, 6}, {2, 7}, {2, 8}, {2, 9}, {3, 4}, {3, 5}, {3, 6}, {3, 7}, {3, 8}, {3, 9}, {4, 6}, {4, 7}, {4, 8}, {4, 10}, {5, 6}, {5, 7}, {5, 8}, {5, 10}, {6, 7}, {6, 8}, {7, 8}}
4	{{1, 4}, {1, 5}, {1, 6}, {1, 7}, {1, 8}, {1, 9}, {2, 4}, {2, 5}, {2, 6}, {2, 7}, {2, 8}, {2, 9}, {3, 4}, {3, 5}, {3, 6}, {3, 7}, {3, 8}, {3, 9}, {4, 7}, {4, 8}, {4, 10}, {5, 7}, {5, 8}, {5, 10}, {6, 7}, {6, 8}, {6, 10}, {7, 8}}
5	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 4}, {2, 5}, {2, 6}, {2, 7}, {2, 8}, {2, 9}, {3, 4}, {3, 5}, {3, 6}, {3, 7}, {3, 8}, {3, 9}, {4, 6}, {4, 7}, {4, 8}, {4, 10}, {5, 6}, {5, 7}, {5, 8}, {5, 10}, {6, 8}, {6, 11}, {7, 8}, {7, 11}}
6	{{1, 6}, {1, 7}, {1, 8}, {1, 9}, {1, 10}, {2, 5}, {2, 7}, {2, 8}, {2, 9}, {2, 11}, {3, 4}, {3, 5}, {3, 6}, {3, 7}, {3, 8}, {3, 9}, {4, 7}, {4, 8}, {4, 10}, {4, 11}, {5, 6}, {5, 7}, {5, 8}, {5, 10}, {6, 7}, {6, 8}, {6, 11}, {7, 8}}
7	{{1, 6}, {1, 7}, {1, 8}, {1, 9}, {1, 10}, {2, 4}, {2, 5}, {2, 8}, {2, 9}, {2, 11}, {3, 4}, {3, 5}, {3, 6}, {3, 7}, {3, 8}, {3, 9}, {4, 6}, {4, 7}, {4, 8}, {4, 10}, {5, 6}, {5, 7}, {5, 8}, {5, 10}, {6, 8}, {6, 11}, {7, 8}, {7, 11}}
8	{{1, 6}, {1, 7}, {1, 8}, {1, 9}, {1, 10}, {2, 4}, {2, 5}, {2, 6}, {2, 7}, {2, 8}, {2, 9}, {3, 4}, {3, 5}, {3, 6}, {3, 7}, {3, 8}, {3, 9}, {4, 6}, {4, 7}, {4, 8}, {4, 10}, {5, 6}, {5, 7}, {5, 8}, {5, 10}, {6, 11}, {7, 11}, {8, 11}}
9	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 5}, {2, 7}, {2, 8}, {2, 9}, {2, 12}, {3, 4}, {3, 5}, {3, 6}, {3, 7}, {3, 8}, {3, 9}, {4, 7}, {4, 8}, {4, 10}, {4, 12}, {5, 6}, {5, 7}, {5, 8}, {5, 10}, {6, 8}, {6, 11}, {6, 12}, {7, 8}, {7, 11}}
10	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 5}, {2, 6}, {2, 7}, {2, 9}, {2, 12}, {3, 4}, {3, 5}, {3, 6}, {3, 7}, {3, 8}, {3, 9}, {4, 6}, {4, 7}, {4, 10}, {4, 12}, {5, 6}, {5, 7}, {5, 8}, {5, 10}, {6, 8}, {6, 11}, {7, 8}, {7, 11}, {8, 12}}
11	{{1, 6}, {1, 7}, {1, 8}, {1, 9}, {1, 10}, {2, 5}, {2, 7}, {2, 8}, {2, 9}, {2, 11}, {3, 4}, {3, 7}, {3, 8}, {3, 9}, {3, 12}, {4, 7}, {4, 8}, {4, 10}, {4, 11}, {5, 7}, {5, 8}, {5, 10}, {5, 12}, {6, 7}, {6, 8}, {6, 11}, {6, 12}, {7, 8}}
12	{{1, 6}, {1, 7}, {1, 8}, {1, 9}, {1, 10}, {2, 5}, {2, 7}, {2, 8}, {2, 9}, {2, 11}, {3, 4}, {3, 6}, {3, 8}, {3, 9}, {3, 12}, {4, 7}, {4, 8}, {4, 10}, {4, 11}, {5, 6}, {5, 8}, {5, 10}, {5, 12}, {6, 7}, {6, 8}, {6, 11}, {7, 8}, {7, 12}}
13	{{1, 6}, {1, 7}, {1, 8}, {1, 9}, {1, 10}, {2, 5}, {2, 7}, {2, 8}, {2, 9}, {2, 11}, {3, 4}, {3, 5}, {3, 6}, {3, 9}, {3, 12}, {4, 7}, {4, 8}, {4, 10}, {4, 11}, {5, 6}, {5, 7}, {5, 8}, {5, 10}, {6, 7}, {6, 8}, {6, 11}, {7, 12}, {8, 12}}
14	{{1, 6}, {1, 7}, {1, 8}, {1, 9}, {1, 10}, {2, 4}, {2, 5}, {2, 8}, {2, 9}, {2, 11}, {3, 4}, {3, 5}, {3, 6}, {3, 7}, {3, 8}, {3, 9}, {4, 7}, {4, 10}, {4, 12}, {5, 6}, {5, 7}, {5, 8}, {5, 10}, {6, 11}, {6, 12}, {7, 8}, {7, 11}, {8, 12}}
15	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 8}, {2, 9}, {2, 12}, {2, 13}, {3, 4}, {3, 5}, {3, 6}, {3, 7}, {3, 8}, {3, 9}, {4, 7}, {4, 8}, {4, 10}, {4, 12}, {5, 6}, {5, 8}, {5, 10}, {5, 13}, {6, 8}, {6, 11}, {6, 12}, {7, 8}, {7, 11}, {7, 13}}
16	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 7}, {2, 9}, {2, 12}, {2, 13}, {3, 4}, {3, 5}, {3, 6}, {3, 7}, {3, 8}, {3, 9}, {4, 7}, {4, 8}, {4, 10}, {4, 12}, {5, 6}, {5, 7}, {5, 10}, {5, 13}, {6, 8}, {6, 11}, {6, 12}, {7, 8}, {7, 11}, {8, 13}}

TABLE 1. Edges Lists for graphs in K_8 family

n	Edges of C_n
17	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 5}, {2, 7}, {2, 8}, {2, 9}, {2, 12}, {3, 4}, {3, 6}, {3, 8}, {3, 9}, {3, 13}, {4, 7}, {4, 8}, {4, 10}, {4, 12}, {5, 6}, {5, 8}, {5, 10}, {5, 13}, {6, 8}, {6, 11}, {6, 12}, {7, 8}, {7, 11}, {7, 13}}
18	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 5}, {2, 7}, {2, 8}, {2, 9}, {2, 12}, {3, 4}, {3, 6}, {3, 7}, {3, 9}, {3, 13}, {4, 7}, {4, 8}, {4, 10}, {4, 12}, {5, 6}, {5, 7}, {5, 10}, {5, 13}, {6, 8}, {6, 11}, {6, 12}, {7, 8}, {7, 11}, {8, 13}}
19	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 5}, {2, 6}, {2, 7}, {2, 9}, {2, 12}, {3, 4}, {3, 6}, {3, 7}, {3, 9}, {3, 13}, {4, 6}, {4, 7}, {4, 10}, {4, 12}, {5, 6}, {5, 7}, {5, 10}, {5, 13}, {6, 8}, {6, 11}, {7, 8}, {7, 11}, {8, 12}, {8, 13}}
20	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 5}, {2, 6}, {2, 7}, {2, 9}, {2, 12}, {3, 4}, {3, 5}, {3, 7}, {3, 9}, {3, 13}, {4, 6}, {4, 7}, {4, 10}, {4, 12}, {5, 6}, {5, 7}, {5, 8}, {5, 10}, {6, 11}, {6, 13}, {7, 8}, {7, 11}, {8, 12}, {8, 13}}
21	{{1, 6}, {1, 7}, {1, 8}, {1, 9}, {1, 10}, {2, 5}, {2, 7}, {2, 8}, {2, 9}, {2, 11}, {3, 4}, {3, 6}, {3, 8}, {3, 9}, {3, 12}, {4, 7}, {4, 8}, {4, 10}, {4, 11}, {5, 6}, {5, 8}, {5, 10}, {5, 12}, {6, 11}, {6, 13}, {7, 12}, {7, 13}, {8, 13}}
22	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 8}, {2, 9}, {2, 12}, {2, 13}, {3, 5}, {3, 6}, {3, 8}, {3, 9}, {3, 14}, {4, 8}, {4, 10}, {4, 12}, {4, 14}, {5, 6}, {5, 8}, {5, 10}, {5, 13}, {6, 8}, {6, 11}, {6, 12}, {7, 8}, {7, 11}, {7, 13}, {7, 14}}
23	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 8}, {2, 9}, {2, 12}, {2, 13}, {3, 5}, {3, 6}, {3, 7}, {3, 9}, {3, 14}, {4, 7}, {4, 10}, {4, 12}, {4, 14}, {5, 6}, {5, 8}, {5, 10}, {5, 13}, {6, 8}, {6, 11}, {6, 12}, {7, 8}, {7, 11}, {7, 13}, {8, 14}}
24	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 8}, {2, 9}, {2, 12}, {2, 13}, {3, 4}, {3, 5}, {3, 6}, {3, 7}, {3, 8}, {3, 9}, {4, 10}, {4, 12}, {4, 14}, {5, 6}, {5, 8}, {5, 10}, {5, 13}, {6, 8}, {6, 11}, {6, 12}, {7, 11}, {7, 13}, {7, 14}, {8, 14}}
25	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 7}, {2, 9}, {2, 12}, {2, 13}, {3, 4}, {3, 6}, {3, 8}, {3, 9}, {3, 14}, {4, 7}, {4, 8}, {4, 10}, {4, 12}, {5, 6}, {5, 10}, {5, 13}, {5, 14}, {6, 8}, {6, 11}, {6, 12}, {7, 8}, {7, 11}, {7, 14}, {8, 13}}
26	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 7}, {2, 9}, {2, 12}, {2, 13}, {3, 4}, {3, 5}, {3, 6}, {3, 9}, {3, 14}, {4, 7}, {4, 8}, {4, 10}, {4, 12}, {5, 6}, {5, 7}, {5, 10}, {5, 13}, {6, 8}, {6, 11}, {6, 12}, {7, 11}, {7, 14}, {8, 13}, {8, 14}}
27	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 5}, {2, 6}, {2, 7}, {2, 9}, {2, 12}, {3, 4}, {3, 5}, {3, 7}, {3, 9}, {3, 13}, {4, 6}, {4, 7}, {4, 10}, {4, 12}, {5, 6}, {5, 10}, {5, 14}, {6, 11}, {6, 13}, {7, 11}, {7, 14}, {8, 12}, {8, 13}, {8, 14}}
28	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 8}, {2, 9}, {2, 12}, {2, 13}, {3, 8}, {3, 9}, {3, 14}, {3, 15}, {4, 8}, {4, 10}, {4, 12}, {4, 14}, {5, 8}, {5, 10}, {5, 13}, {5, 15}, {6, 8}, {6, 11}, {6, 12}, {6, 15}, {7, 8}, {7, 11}, {7, 13}, {7, 14}}
29	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 8}, {2, 9}, {2, 12}, {2, 13}, {3, 6}, {3, 9}, {3, 14}, {3, 15}, {4, 8}, {4, 10}, {4, 12}, {4, 14}, {5, 6}, {5, 10}, {5, 13}, {5, 15}, {6, 8}, {6, 11}, {6, 12}, {7, 8}, {7, 11}, {7, 13}, {7, 14}, {8, 15}}
30	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 8}, {2, 9}, {2, 12}, {2, 13}, {3, 5}, {3, 6}, {3, 7}, {3, 9}, {3, 14}, {4, 7}, {4, 10}, {4, 12}, {4, 14}, {5, 10}, {5, 13}, {5, 15}, {6, 11}, {6, 12}, {6, 15}, {7, 8}, {7, 11}, {7, 13}, {8, 14}, {8, 15}}
31	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 7}, {2, 9}, {2, 12}, {2, 13}, {3, 4}, {3, 9}, {3, 14}, {3, 15}, {4, 7}, {4, 8}, {4, 10}, {4, 12}, {5, 6}, {5, 10}, {5, 13}, {5, 14}, {6, 11}, {6, 12}, {6, 15}, {7, 8}, {7, 11}, {7, 14}, {8, 13}, {8, 15}}
32	{{1, 8}, {1, 9}, {1, 10}, {1, 11}, {2, 7}, {2, 9}, {2, 12}, {2, 13}, {3, 4}, {3, 9}, {3, 14}, {3, 15}, {4, 10}, {4, 12}, {4, 16}, {5, 6}, {5, 10}, {5, 13}, {5, 14}, {6, 11}, {6, 12}, {6, 15}, {7, 11}, {7, 14}, {7, 16}, {8, 13}, {8, 15}, {8, 16}}

TABLE 2. Edges Lists for graphs in K_8 family (continued)

n	Edges deleted/contracted	$K_{3^2,1^2}$ cousin
5	$\{\{4,6\},\{4,8\},\{6,8\}\}$	4
6	$\{\{4,7\},\{4,8\},\{7,8\}\}$	4
7	$\{\{4,6\},\{4,8\},\{6,8\}\}$	6
8	$\{\{4,6\},\{4,7\}\}$ / $\{8,11\}$	2
9	$\{\{4,7\},\{4,8\},\{7,8\}\}$	8
10	$\{\{5,6\},\{5,8\},\{6,8\}\}$	9
11	$\{\{1,7\},\{1,8\},\{7,8\}\}$	11
12	$\{\{4,7\},\{4,8\},\{7,8\}\}$	9
13	$\{\{5,6\},\{5,7\},\{6,7\}\}$	14
14	$\{\{5,7\},\{5,8\},\{7,8\}\}$	16
15	$\{\{4,7\},\{4,8\},\{7,8\}\}$	17
16	$\{\{4,7\},\{4,8\},\{7,8\}\}$	18
17	$\{\{4,7\},\{4,8\},\{7,8\}\}$	18
18	$\{\{4,7\},\{4,8\},\{7,8\}\}$	19
19	/ $\{\{1,10\},\{2,12\},\{3,13\}\}$	3
20	$\{\{5,7\},\{5,7\},\{7,8\}\}$	23
21	$\{4,7\}$ / $\{\{2,11\},\{8,13\}\}$	5
22	$\{\{5,6\},\{5,8\},\{6,8\}\}$	29
23	$\{\{5,6\},\{5,8\},\{6,8\}\}$	29
24	$\{\{5,6\},\{5,8\},\{6,8\}\}$	29
25	$\{\{4,7\},\{4,8\},\{7,8\}\}$	30
26	$\{5,7\}$ / $\{\{2,13\},\{3,14\}\}$	10
27	/ $\{\{1,10\},\{2,12\},\{7,14\}\}$	7
28	/ $\{\{1,10\},\{2,12\},\{3,15\}\}$	13
29	/ $\{\{1,10\},\{2,13\},\{3,14\}\}$	14
30	/ $\{\{1,10\},\{3,14\},\{6,15\}\}$	10
31	$\{\{4,7\},\{4,8\},\{7,8\}\}$	42
32	/ $\{\{1,10\},\{2,13\},\{7,16\}\}$	24

TABLE 3. Identifying a minor of $C_n - 9$ in the $K_{3^2,1^2}$ family

DEPARTMENT OF MATHEMATICS AND STATISTICS, CALIFORNIA STATE UNIVERSITY, CHICO, CHICO, CA 95929-0525

E-mail address: TMattman@CSUChico.edu

DEPARTMENT OF MATHEMATICS, UC, RIVERSIDE, 900 UNIVERSITY AVENUE, RIVERSIDE, CA 92521

E-mail address: mpierce@math.ucr.edu

n	Edge e	Apex Sets
5	$\{1,8\}$	$C_5 - e$: 2, 3, 4; C_5/e: 1, 2, 3
	$\{1,9\}$	$C_5 - e$: 2, 4, 5; C_5/e: 1, 4, 5
	$\{2,4\}$	$C_5 - e$: 1, 3, 5; C_5/e: 1, 2, 3
	$\{2,8\}$	$C_5 - e$: 1, 3, 4; C_5/e: 1, 2, 3
	$\{2,9\}$	$C_5 - e$: 1, 4, 5; C_5/e: 2, 4, 5
6	$\{1,6\}$	$C_6 - e$: 2, 3, 4; C_6/e: 1, 2, 3
	$\{1,7\}$	$C_6 - e$: 2, 3, 4; C_6/e: 1, 2, 3
	$\{1,9\}$	$C_6 - e$: 2, 4, 5; C_6/e: 1, 4, 5
	$\{3,5\}$	$C_6 - e$: 1, 2, 4; C_6/e: 1, 2, 3
	$\{3,7\}$	$C_6 - e$: 1, 2, 4; C_6/e: 1, 2, 3
	$\{3,9\}$	$C_6 - e$: 1, 4, 5; C_6/e: 3, 4, 5
	$\{7,8\}$	$C_6 - e$: 1, 2, 3; C_6/e: 1, 2, 3
7	$\{1,6\}$	$C_7 - e$: 2, 3, 4; C_7/e: 1, 2, 3
	$\{1,8\}$	$C_7 - e$: 2, 3, 4; C_7/e: 1, 2, 3
	$\{1,9\}$	$C_7 - e$: 2, 4, 5; C_7/e: 1, 4, 5
	$\{1,10\}$	$C_7 - e$: 2, 3, 4; C_7/e: 1, 2, 3
	$\{3,4\}$	$C_7 - e$: 1, 2, 5; C_7/e: 1, 2, 3
	$\{3,8\}$	$C_7 - e$: 1, 2, 4; C_7/e: 1, 2, 3
	$\{3,9\}$	$C_7 - e$: 1, 4, 5; C_7/e: 3, 4, 5
	$\{4,6\}$	$C_7 - e$: 1, 2, 3; C_7/e: 1, 2, 3
	$\{4,8\}$	$C_7 - e$: 1, 2, 3; C_7/e: 1, 2, 3
	$\{4,10\}$	$C_7 - e$: 1, 2, 3; C_7/e: 2, 3, 4
8	$\{1,6\}$	$C_8 - e$: 2, 3, 4; C_8/e: 1, 2, 3
	$\{1,9\}$	$C_8 - e$: 2, 4, 6; C_8/e: 1, 4, 6
	$\{2,4\}$	$C_8 - e$: 1, 3, 6; C_8/e: 1, 2, 3
	$\{2,6\}$	$C_8 - e$: 1, 3, 4; C_8/e: 1, 2, 3
	$\{2,9\}$	$C_8 - e$: 1, 4, 6; C_8/e: 2, 4, 6
	$\{6,11\}$	$C_8 - e$: 1, 2, 3; C_8/e: 1, 2, 3
9	$\{1,8\}$	$C_9 - e$: 2, 3, 4; C_9/e: 1, 2, 3
	$\{1,9\}$	$C_9 - e$: 2, 4, 5; C_9/e: 1, 4, 5
	$\{2,5\}$	$C_9 - e$: 1, 3, 4; C_9/e: 1, 2, 3
	$\{2,8\}$	$C_9 - e$: 1, 3, 4; C_9/e: 1, 2, 3
	$\{2,9\}$	$C_9 - e$: 1, 4, 5; C_9/e: 2, 4, 5
	$\{2,12\}$	$C_9 - e$: 1, 3, 4; C_9/e: 1, 2, 3
	$\{3,5\}$	$C_9 - e$: 1, 2, 4; C_9/e: 1, 2, 3
	$\{3,8\}$	$C_9 - e$: 1, 2, 4; C_9/e: 1, 2, 3
	$\{3,9\}$	$C_9 - e$: 1, 4, 5; C_9/e: 3, 4, 5

TABLE 4. Every proper minor of C_n is 3-apex.

n	Edge e	Apex Sets
10	$\{1,8\}$	$C_{10}-e$: 2, 3, 4; C_{10}/e: 1, 2, 3
	$\{1,9\}$	$C_{10}-e$: 2, 4, 5; C_{10}/e: 1, 4, 5
	$\{1,11\}$	$C_{10}-e$: 2, 3, 4; C_{10}/e: 1, 2, 3
	$\{2,5\}$	$C_{10}-e$: 1, 3, 4; C_{10}/e: 1, 2, 3
	$\{2,6\}$	$C_{10}-e$: 1, 3, 4; C_{10}/e: 1, 2, 3
	$\{2,9\}$	$C_{10}-e$: 1, 4, 5; C_{10}/e: 2, 4, 5
	$\{2,12\}$	$C_{10}-e$: 1, 3, 4; C_{10}/e: 1, 2, 3
	$\{3,5\}$	$C_{10}-e$: 1, 2, 4; C_{10}/e: 1, 2, 3
	$\{3,6\}$	$C_{10}-e$: 1, 2, 4; C_{10}/e: 1, 2, 3
	$\{3,8\}$	$C_{10}-e$: 1, 2, 4; C_{10}/e: 1, 2, 3
	$\{3,9\}$	$C_{10}-e$: 1, 4, 5; C_{10}/e: 3, 4, 5
	$\{6,8\}$	$C_{10}-e$: 1, 2, 3; C_{10}/e: 1, 2, 3
	$\{6,11\}$	$C_{10}-e$: 1, 2, 3; C_{10}/e: 2, 3, 4
	$\{8,12\}$	$C_{10}-e$: 1, 2, 3; C_{10}/e: 1, 3, 5
11	$\{1,6\}$	$C_{11}-e$: 2, 3, 4; C_{11}/e: 1, 2, 3
	$\{1,7\}$	$C_{11}-e$: 2, 3, 4; C_{11}/e: 1, 2, 3
	$\{1,9\}$	$C_{11}-e$: 2, 4, 5; C_{11}/e: 1, 4, 5
	$\{7,8\}$	$C_{11}-e$: 1, 2, 3; C_{11}/e: 1, 2, 3
12	$\{1,6\}$	$C_{12}-e$: 2, 3, 4; C_{12}/e: 1, 2, 3
	$\{1,8\}$	$C_{12}-e$: 2, 3, 4; C_{12}/e: 1, 2, 3
	$\{1,9\}$	$C_{12}-e$: 2, 4, 5; C_{12}/e: 1, 4, 5
	$\{2,5\}$	$C_{12}-e$: 1, 3, 4; C_{12}/e: 1, 2, 3
	$\{2,7\}$	$C_{12}-e$: 1, 3, 4; C_{12}/e: 1, 2, 3
	$\{2,8\}$	$C_{12}-e$: 1, 3, 4; C_{12}/e: 1, 2, 3
	$\{2,9\}$	$C_{12}-e$: 1, 4, 5; C_{12}/e: 2, 4, 5
	$\{2,11\}$	$C_{12}-e$: 1, 3, 4; C_{12}/e: 1, 2, 3
	$\{6,7\}$	$C_{12}-e$: 1, 2, 3; C_{12}/e: 1, 2, 3
	$\{6,8\}$	$C_{12}-e$: 1, 2, 3; C_{12}/e: 1, 2, 3
	$\{6,11\}$	$C_{12}-e$: 1, 2, 3; C_{12}/e: 1, 3, 5
13	$\{1,6\}$	$C_{13}-e$: 2, 3, 4; C_{13}/e: 1, 2, 3
	$\{1,7\}$	$C_{13}-e$: 2, 3, 4; C_{13}/e: 1, 2, 3
	$\{1,9\}$	$C_{13}-e$: 2, 4, 5; C_{13}/e: 1, 4, 5
	$\{1,10\}$	$C_{13}-e$: 2, 3, 4; C_{13}/e: 1, 2, 3
	$\{3,4\}$	$C_{13}-e$: 1, 2, 5; C_{13}/e: 1, 2, 3
	$\{3,5\}$	$C_{13}-e$: 1, 2, 4; C_{13}/e: 1, 2, 3
	$\{3,9\}$	$C_{13}-e$: 1, 4, 5; C_{13}/e: 3, 4, 5
	$\{3,12\}$	$C_{13}-e$: 1, 2, 4; C_{13}/e: 1, 2, 3
	$\{4,7\}$	$C_{13}-e$: 1, 2, 3; C_{13}/e: 1, 2, 3
	$\{4,10\}$	$C_{13}-e$: 1, 2, 3; C_{13}/e: 2, 3, 4
	$\{5,6\}$	$C_{13}-e$: 1, 2, 3; C_{13}/e: 1, 2, 3
	$\{5,7\}$	$C_{13}-e$: 1, 2, 3; C_{13}/e: 1, 2, 3
	$\{5,10\}$	$C_{13}-e$: 1, 2, 3; C_{13}/e: 2, 3, 5
	$\{7,12\}$	$C_{13}-e$: 1, 2, 3; C_{13}/e: 1, 2, 4

TABLE 5. Every proper minor of C_n is 3-apex (continued)

n	Edge e	Apex Sets
14	$\{1,6\}$	$C_{14}-e$: 2, 3, 4; C_{14}/e: 1, 2, 3
	$\{1,7\}$	$C_{14}-e$: 2, 3, 4; C_{14}/e: 1, 2, 3
	$\{1,9\}$	$C_{14}-e$: 2, 4, 5; C_{14}/e: 1, 4, 5
	$\{3,5\}$	$C_{14}-e$: 1, 2, 4; C_{14}/e: 1, 2, 3
	$\{3,8\}$	$C_{14}-e$: 1, 2, 4; C_{14}/e: 1, 2, 3
	$\{3,9\}$	$C_{14}-e$: 1, 4, 5; C_{14}/e: 3, 4, 5
15	$\{1,8\}$	$C_{15}-e$: 2, 3, 4; C_{15}/e: 1, 2, 3
	$\{1,9\}$	$C_{15}-e$: 2, 4, 5; C_{15}/e: 1, 4, 5
	$\{1,10\}$	$C_{15}-e$: 2, 3, 4; C_{15}/e: 1, 2, 3
	$\{3,4\}$	$C_{15}-e$: 1, 2, 5; C_{15}/e: 1, 2, 3
	$\{3,8\}$	$C_{15}-e$: 1, 2, 4; C_{15}/e: 1, 2, 3
	$\{3,9\}$	$C_{15}-e$: 1, 4, 5; C_{15}/e: 3, 4, 5
	$\{4,7\}$	$C_{15}-e$: 1, 2, 3; C_{15}/e: 1, 2, 3
	$\{4,8\}$	$C_{15}-e$: 1, 2, 3; C_{15}/e: 1, 2, 3
	$\{4,10\}$	$C_{15}-e$: 1, 2, 3; C_{15}/e: 2, 3, 4
16	$\{1,8\}$	$C_{16}-e$: 2, 3, 4; C_{16}/e: 1, 2, 3
	$\{1,9\}$	$C_{16}-e$: 2, 4, 5; C_{16}/e: 1, 4, 5
	$\{1,10\}$	$C_{16}-e$: 2, 3, 4; C_{16}/e: 1, 2, 3
	$\{1,11\}$	$C_{16}-e$: 2, 3, 4; C_{16}/e: 1, 2, 3
	$\{3,4\}$	$C_{16}-e$: 1, 2, 5; C_{16}/e: 1, 2, 3
	$\{3,5\}$	$C_{16}-e$: 1, 2, 4; C_{16}/e: 1, 2, 3
	$\{3,7\}$	$C_{16}-e$: 1, 2, 4; C_{16}/e: 1, 2, 3
	$\{3,9\}$	$C_{16}-e$: 1, 4, 5; C_{16}/e: 3, 4, 5
	$\{4,7\}$	$C_{16}-e$: 1, 2, 3; C_{16}/e: 1, 2, 3
	$\{4,10\}$	$C_{16}-e$: 1, 2, 3; C_{16}/e: 2, 3, 4
	$\{5,6\}$	$C_{16}-e$: 1, 2, 3; C_{16}/e: 1, 2, 3
	$\{5,7\}$	$C_{16}-e$: 1, 2, 3; C_{16}/e: 1, 2, 3
	$\{5,10\}$	$C_{16}-e$: 1, 2, 3; C_{16}/e: 2, 3, 5
	$\{5,13\}$	$C_{16}-e$: 1, 2, 3; C_{16}/e: 1, 3, 4
	$\{7,8\}$	$C_{16}-e$: 1, 2, 3; C_{16}/e: 1, 2, 3
	$\{7,11\}$	$C_{16}-e$: 1, 2, 3; C_{16}/e: 2, 3, 4
17	$\{1,8\}$	$C_{17}-e$: 2, 3, 4; C_{17}/e: 1, 2, 3
	$\{1,9\}$	$C_{17}-e$: 2, 4, 5; C_{17}/e: 1, 4, 5
	$\{2,5\}$	$C_{17}-e$: 1, 3, 4; C_{17}/e: 1, 2, 3
	$\{2,8\}$	$C_{17}-e$: 1, 3, 4; C_{17}/e: 1, 2, 3
	$\{2,9\}$	$C_{17}-e$: 1, 4, 5; C_{17}/e: 2, 4, 5
	$\{2,12\}$	$C_{17}-e$: 1, 3, 4; C_{17}/e: 1, 2, 3

TABLE 6. Every proper minor of C_n is 3-apex (continued)

n	Edge e	Apex Sets
18	$\{1,8\}$	$C_{18}-e$: 2, 3, 4; C_{18}/e: 1, 2, 3
	$\{1,9\}$	$C_{18}-e$: 2, 4, 5; C_{18}/e: 1, 4, 5
	$\{1,11\}$	$C_{18}-e$: 2, 3, 4; C_{18}/e: 1, 2, 3
	$\{2,5\}$	$C_{18}-e$: 1, 3, 4; C_{18}/e: 1, 2, 3
	$\{2,7\}$	$C_{18}-e$: 1, 3, 4; C_{18}/e: 1, 2, 3
	$\{2,8\}$	$C_{18}-e$: 1, 3, 4; C_{18}/e: 1, 2, 3
	$\{2,9\}$	$C_{18}-e$: 1, 4, 5; C_{18}/e: 2, 4, 5
	$\{2,12\}$	$C_{18}-e$: 1, 3, 4; C_{18}/e: 1, 2, 3
	$\{3,6\}$	$C_{18}-e$: 1, 2, 4; C_{18}/e: 1, 2, 3
	$\{3,7\}$	$C_{18}-e$: 1, 2, 4; C_{18}/e: 1, 2, 3
	$\{3,9\}$	$C_{18}-e$: 1, 4, 5; C_{18}/e: 3, 4, 5
	$\{3,13\}$	$C_{18}-e$: 1, 2, 4; C_{18}/e: 1, 2, 3
	$\{6,8\}$	$C_{18}-e$: 1, 2, 3; C_{18}/e: 1, 2, 3
	$\{6,11\}$	$C_{18}-e$: 1, 2, 3; C_{18}/e: 2, 3, 4
	$\{6,12\}$	$C_{18}-e$: 1, 2, 3; C_{18}/e: 1, 3, 5
	$\{7,8\}$	$C_{18}-e$: 1, 2, 3; C_{18}/e: 1, 2, 3
	$\{7,11\}$	$C_{18}-e$: 1, 2, 3; C_{18}/e: 2, 3, 4
	$\{8,13\}$	$C_{18}-e$: 1, 2, 3; C_{18}/e: 1, 2, 4
19	$\{1,8\}$	$C_{19}-e$: 2, 3, 4; C_{19}/e: 1, 2, 3
	$\{1,9\}$	$C_{19}-e$: 2, 4, 5; C_{19}/e: 1, 4, 5
	$\{1,11\}$	$C_{19}-e$: 2, 3, 4; C_{19}/e: 1, 2, 3
	$\{2,5\}$	$C_{19}-e$: 1, 3, 4; C_{19}/e: 1, 2, 3
	$\{2,6\}$	$C_{19}-e$: 1, 3, 4; C_{19}/e: 1, 2, 3
	$\{2,9\}$	$C_{19}-e$: 1, 4, 5; C_{19}/e: 2, 4, 5
	$\{2,12\}$	$C_{19}-e$: 1, 3, 4; C_{19}/e: 1, 2, 3
	$\{6,8\}$	$C_{19}-e$: 1, 2, 3; C_{19}/e: 1, 2, 3
	$\{6,11\}$	$C_{19}-e$: 1, 2, 3; C_{19}/e: 2, 3, 4
	$\{8,12\}$	$C_{19}-e$: 1, 2, 3; C_{19}/e: 1, 3, 5
20	$\{1,8\}$	$C_{20}-e$: 2, 3, 4; C_{20}/e: 1, 2, 3
	$\{1,9\}$	$C_{20}-e$: 2, 4, 5; C_{20}/e: 1, 4, 5
	$\{1,10\}$	$C_{20}-e$: 2, 3, 4; C_{20}/e: 1, 2, 3
	$\{2,5\}$	$C_{20}-e$: 1, 3, 4; C_{20}/e: 1, 2, 3
	$\{2,6\}$	$C_{20}-e$: 1, 3, 4; C_{20}/e: 1, 2, 3
	$\{2,7\}$	$C_{20}-e$: 1, 3, 4; C_{20}/e: 1, 2, 3
	$\{2,9\}$	$C_{20}-e$: 1, 4, 5; C_{20}/e: 2, 4, 5
	$\{2,12\}$	$C_{20}-e$: 1, 3, 4; C_{20}/e: 1, 2, 3
	$\{4,6\}$	$C_{20}-e$: 1, 2, 3; C_{20}/e: 1, 2, 3
	$\{4,7\}$	$C_{20}-e$: 1, 2, 3; C_{20}/e: 1, 2, 3
	$\{4,10\}$	$C_{20}-e$: 1, 2, 3; C_{20}/e: 2, 3, 4
	$\{4,12\}$	$C_{20}-e$: 1, 2, 3; C_{20}/e: 1, 3, 4
	$\{5,7\}$	$C_{20}-e$: 1, 2, 3; C_{20}/e: 1, 2, 3
	$\{5,8\}$	$C_{20}-e$: 1, 2, 3; C_{20}/e: 1, 2, 3
	$\{5,10\}$	$C_{20}-e$: 1, 2, 3; C_{20}/e: 2, 3, 5
	$\{8,12\}$	$C_{20}-e$: 1, 2, 3; C_{20}/e: 1, 3, 5

TABLE 7. Every proper minor of C_n is 3-apex (continued)

n	Edge e	Apex Sets
21	$\{1,6\}$	$C_{21}-e$: 2, 3, 4; C_{21}/e: 1, 2, 3
	$\{1,8\}$	$C_{21}-e$: 2, 3, 4; C_{21}/e: 1, 2, 3
	$\{1,9\}$	$C_{21}-e$: 2, 4, 6; C_{21}/e: 1, 4, 6
	$\{2,5\}$	$C_{21}-e$: 1, 3, 6; C_{21}/e: 1, 2, 3
	$\{2,7\}$	$C_{21}-e$: 1, 3, 4; C_{21}/e: 1, 2, 3
	$\{2,8\}$	$C_{21}-e$: 1, 3, 4; C_{21}/e: 1, 2, 3
	$\{2,9\}$	$C_{21}-e$: 1, 4, 6; C_{21}/e: 2, 4, 6
	$\{2,11\}$	$C_{21}-e$: 1, 3, 4; C_{21}/e: 1, 2, 3
	$\{6,11\}$	$C_{21}-e$: 1, 2, 3; C_{21}/e: 1, 3, 6
	$\{6,13\}$	$C_{21}-e$: 1, 2, 3; C_{21}/e: 1, 2, 3
	$\{8,13\}$	$C_{21}-e$: 1, 2, 3; C_{21}/e: 1, 2, 3
22	$\{1,8\}$	$C_{22}-e$: 2, 3, 4; C_{22}/e: 1, 2, 3
	$\{1,9\}$	$C_{22}-e$: 2, 4, 5; C_{22}/e: 1, 4, 5
	$\{3,5\}$	$C_{22}-e$: 1, 2, 4; C_{22}/e: 1, 2, 3
	$\{3,8\}$	$C_{22}-e$: 1, 2, 4; C_{22}/e: 1, 2, 3
	$\{3,9\}$	$C_{22}-e$: 1, 4, 5; C_{22}/e: 3, 4, 5
23	$\{1,8\}$	$C_{23}-e$: 2, 3, 4; C_{23}/e: 1, 2, 3
	$\{1,9\}$	$C_{23}-e$: 2, 4, 5; C_{23}/e: 1, 4, 5
	$\{1,10\}$	$C_{23}-e$: 2, 3, 4; C_{23}/e: 1, 2, 3
	$\{1,11\}$	$C_{23}-e$: 2, 3, 4; C_{23}/e: 1, 2, 3
	$\{3,5\}$	$C_{23}-e$: 1, 2, 4; C_{23}/e: 1, 2, 3
	$\{3,7\}$	$C_{23}-e$: 1, 2, 4; C_{23}/e: 1, 2, 3
	$\{3,9\}$	$C_{23}-e$: 1, 4, 5; C_{23}/e: 3, 4, 5
	$\{3,14\}$	$C_{23}-e$: 1, 2, 4; C_{23}/e: 1, 2, 3
	$\{4,7\}$	$C_{23}-e$: 1, 2, 3; C_{23}/e: 1, 2, 3
	$\{4,10\}$	$C_{23}-e$: 1, 2, 3; C_{23}/e: 2, 3, 4
	$\{4,14\}$	$C_{23}-e$: 1, 2, 3; C_{23}/e: 1, 2, 4
	$\{5,6\}$	$C_{23}-e$: 1, 2, 3; C_{23}/e: 1, 2, 3
	$\{5,8\}$	$C_{23}-e$: 1, 2, 3; C_{23}/e: 1, 2, 3
	$\{5,10\}$	$C_{23}-e$: 1, 2, 3; C_{23}/e: 2, 3, 5
	$\{5,13\}$	$C_{23}-e$: 1, 2, 3; C_{23}/e: 1, 3, 4
	$\{7,8\}$	$C_{23}-e$: 1, 2, 3; C_{23}/e: 1, 2, 3
	$\{7,11\}$	$C_{23}-e$: 1, 2, 3; C_{23}/e: 2, 3, 4
	$\{8,14\}$	$C_{23}-e$: 1, 2, 3; C_{23}/e: 1, 2, 5
24	$\{1,8\}$	$C_{24}-e$: 2, 3, 4; C_{24}/e: 1, 2, 3
	$\{1,9\}$	$C_{24}-e$: 2, 4, 5; C_{24}/e: 1, 4, 5
	$\{1,10\}$	$C_{24}-e$: 2, 3, 4; C_{24}/e: 1, 2, 3
	$\{3,5\}$	$C_{24}-e$: 1, 2, 4; C_{24}/e: 1, 2, 3
	$\{3,8\}$	$C_{24}-e$: 1, 2, 4; C_{24}/e: 1, 2, 3
	$\{3,9\}$	$C_{24}-e$: 1, 4, 5; C_{24}/e: 3, 4, 5
	$\{5,6\}$	$C_{24}-e$: 1, 2, 4; C_{24}/e: 1, 2, 3
	$\{5,10\}$	$C_{24}-e$: 1, 2, 3; C_{24}/e: 2, 3, 5

TABLE 8. Every proper minor of C_n is 3-apex (continued)

n	Edge e	Apex Sets
25	$\{1,8\}$	$C_{25}-e$: 2, 3, 4; C_{25}/e: 1, 2, 3
	$\{1,9\}$	$C_{25}-e$: 2, 4, 5; C_{25}/e: 1, 4, 5
	$\{1,11\}$	$C_{25}-e$: 2, 3, 4; C_{25}/e: 1, 2, 3
	$\{2,7\}$	$C_{25}-e$: 1, 3, 4; C_{25}/e: 1, 2, 3
	$\{2,9\}$	$C_{25}-e$: 1, 4, 5; C_{25}/e: 2, 4, 5
	$\{2,12\}$	$C_{25}-e$: 1, 3, 4; C_{25}/e: 1, 2, 3
	$\{2,13\}$	$C_{25}-e$: 1, 3, 4; C_{25}/e: 1, 2, 3
	$\{3,4\}$	$C_{25}-e$: 1, 2, 5; C_{25}/e: 1, 2, 3
	$\{3,6\}$	$C_{25}-e$: 1, 2, 4; C_{25}/e: 1, 2, 3
	$\{3,8\}$	$C_{25}-e$: 1, 2, 4; C_{25}/e: 1, 2, 3
	$\{3,9\}$	$C_{25}-e$: 1, 4, 5; C_{25}/e: 3, 4, 5
	$\{3,14\}$	$C_{25}-e$: 1, 2, 4; C_{25}/e: 1, 2, 3
	$\{6,8\}$	$C_{25}-e$: 1, 2, 3; C_{25}/e: 1, 2, 3
	$\{6,11\}$	$C_{25}-e$: 1, 2, 3; C_{25}/e: 2, 3, 4
	$\{6,12\}$	$C_{25}-e$: 1, 2, 3; C_{25}/e: 1, 3, 5
	$\{8,13\}$	$C_{25}-e$: 1, 2, 3; C_{25}/e: 1, 3, 4
26	$\{1,8\}$	$C_{26}-e$: 2, 3, 4; C_{26}/e: 1, 2, 3
	$\{1,9\}$	$C_{26}-e$: 2, 4, 5; C_{26}/e: 1, 4, 5
	$\{1,10\}$	$C_{26}-e$: 2, 3, 4; C_{26}/e: 1, 2, 3
	$\{1,11\}$	$C_{26}-e$: 2, 3, 4; C_{26}/e: 1, 2, 3
	$\{3,4\}$	$C_{26}-e$: 1, 2, 5; C_{26}/e: 1, 2, 3
	$\{3,5\}$	$C_{26}-e$: 1, 2, 4; C_{26}/e: 1, 2, 3
	$\{3,9\}$	$C_{26}-e$: 1, 4, 5; C_{26}/e: 3, 4, 5
	$\{3,14\}$	$C_{26}-e$: 1, 2, 4; C_{26}/e: 1, 2, 3
	$\{4,7\}$	$C_{26}-e$: 1, 2, 3; C_{26}/e: 1, 2, 3
	$\{4,10\}$	$C_{26}-e$: 1, 2, 3; C_{26}/e: 2, 3, 4
	$\{5,6\}$	$C_{26}-e$: 1, 2, 3; C_{26}/e: 1, 2, 3
	$\{5,7\}$	$C_{26}-e$: 1, 2, 3; C_{26}/e: 1, 2, 3
	$\{5,10\}$	$C_{26}-e$: 1, 2, 3; C_{26}/e: 2, 3, 5
	$\{5,13\}$	$C_{26}-e$: 1, 2, 3; C_{26}/e: 1, 3, 4
	$\{7,11\}$	$C_{26}-e$: 1, 2, 3; C_{26}/e: 2, 3, 4
	$\{7,14\}$	$C_{26}-e$: 1, 2, 3; C_{26}/e: 1, 2, 4
27	$\{1,8\}$	$C_{27}-e$: 2, 3, 4; C_{27}/e: 1, 2, 3
	$\{1,9\}$	$C_{27}-e$: 2, 4, 5; C_{27}/e: 1, 4, 5
	$\{2,5\}$	$C_{27}-e$: 1, 3, 4; C_{27}/e: 1, 2, 3
	$\{2,7\}$	$C_{27}-e$: 1, 3, 4; C_{27}/e: 1, 2, 3
	$\{2,9\}$	$C_{27}-e$: 1, 4, 5; C_{27}/e: 2, 4, 5
28	$\{1,8\}$	$C_{28}-e$: 2, 3, 4; C_{28}/e: 1, 2, 3
	$\{1,9\}$	$C_{28}-e$: 2, 4, 5; C_{28}/e: 1, 4, 5

TABLE 9. Every proper minor of C_n is 3-apex. (continued)

n	Edge e	Apex Sets
29	$\{1,8\}$	$C_{29}-e$: 2, 3, 4; C_{29}/e: 1, 2, 3
	$\{1,9\}$	$C_{29}-e$: 2, 4, 5; C_{29}/e: 1, 4, 5
	$\{1,11\}$	$C_{29}-e$: 2, 3, 4; C_{29}/e: 1, 2, 3
	$\{3,6\}$	$C_{29}-e$: 1, 2, 4; C_{29}/e: 1, 2, 3
	$\{3,9\}$	$C_{29}-e$: 1, 4, 5; C_{29}/e: 3, 4, 5
	$\{3,15\}$	$C_{29}-e$: 1, 2, 4; C_{29}/e: 1, 2, 3
	$\{6,8\}$	$C_{29}-e$: 1, 2, 3; C_{29}/e: 1, 2, 3
	$\{6,11\}$	$C_{29}-e$: 1, 2, 3; C_{29}/e: 2, 3, 4
	$\{8,15\}$	$C_{29}-e$: 1, 2, 3; C_{29}/e: 1, 2, 4
30	$\{1,8\}$	$C_{30}-e$: 2, 3, 4; C_{30}/e: 1, 2, 3
	$\{1,9\}$	$C_{30}-e$: 2, 4, 5; C_{30}/e: 1, 4, 5
	$\{1,10\}$	$C_{30}-e$: 2, 3, 4; C_{30}/e: 1, 2, 3
	$\{1,11\}$	$C_{30}-e$: 2, 3, 4; C_{30}/e: 1, 2, 3
	$\{3,7\}$	$C_{30}-e$: 1, 2, 5; C_{30}/e: 1, 2, 3
	$\{3,9\}$	$C_{30}-e$: 1, 4, 5; C_{30}/e: 3, 4, 5
	$\{3,14\}$	$C_{30}-e$: 1, 2, 4; C_{30}/e: 1, 2, 3
	$\{4,7\}$	$C_{30}-e$: 1, 2, 5; C_{30}/e: 1, 2, 3
	$\{4,10\}$	$C_{30}-e$: 1, 2, 3; C_{30}/e: 2, 3, 4
	$\{4,14\}$	$C_{30}-e$: 1, 2, 3; C_{30}/e: 1, 2, 4
	$\{7,11\}$	$C_{30}-e$: 1, 2, 3; C_{30}/e: 2, 3, 4
31	$\{1,8\}$	$C_{31}-e$: 2, 3, 4; C_{31}/e: 1, 2, 3
	$\{1,9\}$	$C_{31}-e$: 2, 4, 5; C_{31}/e: 1, 4, 5
	$\{1,10\}$	$C_{31}-e$: 2, 3, 4; C_{31}/e: 1, 2, 3
	$\{4,7\}$	$C_{31}-e$: 1, 2, 3; C_{31}/e: 1, 2, 3
	$\{4,10\}$	$C_{31}-e$: 1, 2, 3; C_{31}/e: 2, 3, 4
	$\{5,6\}$	$C_{31}-e$: 1, 2, 3; C_{31}/e: 1, 2, 3
	$\{5,10\}$	$C_{31}-e$: 1, 2, 3; C_{31}/e: 2, 3, 5
32	$\{1,8\}$	$C_{32}-e$: 2, 3, 4; C_{32}/e: 1, 2, 3
	$\{1,9\}$	$C_{32}-e$: 2, 4, 5; C_{32}/e: 1, 4, 5

TABLE 10. Every proper minor of C_n is 3-apex (continued)